

Overview

Method Name

High Performance Liquid Chromatography with Ultraviolet Detection (HPLC-UV)

NY State Available

Yes

Specimen

Specimen Type

Varies

Specimen Required

Submit only 1 of the following specimens:

Plasma

Draw blood in a green-top (sodium heparin) tube(s), **plasma gel tube is not acceptable**. Spin down and send 2 mL sodium heparin plasma refrigerated in a plastic vial.

Serum

Draw blood in a plain red-top tube(s), **serum gel tube is not acceptable**. Spin down and send 2 mL of serum refrigerated in a plastic vial.

Specimen Minimum Volume

0.3 mL

Reject Due To

Hemolysis	NA
Lipemia	NA
Icterus	NA
Other	NA

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Refrigerated (preferred)	14 days	
	Frozen	180 days	
	Ambient	72 hours	

Clinical and Interpretive

Reference Values

Units: ng/mL

Plasma insulin concentrations have been shown to increase only when plasma glipizide concentrations exceeded 200 ng/mL.

Toxic range has not been established.

Performance

PDF Report

No

Day(s) and Time(s) Test Performed

Monday through Sunday

Analytic Time

7 days

Maximum Laboratory Time

9 - 10 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Medtox Laboratories, Inc.

Fees and Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

CPT Code Information

80375

G0480 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
FGLIP	Glipizide (Glucotrol)	10539-5

Result ID	Test Result Name	Result LOINC Value
Z1180	Glipizide	10539-5